

AD-A048 061

OFFICE OF NAVAL RESEARCH LONDON (ENGLAND)

F/G 20/8

ACOUSTIC DETECTION OF NEUTRINO INTERACTIONS IN THE OCEAN: THE 1--ETC(U).

NOV 77 A ROBERTS

UNCLASSIFIED

ONRL-C-15-77

NL

| OF |  
AD  
A048061



END  
DATE  
FILMED  
1-78  
DDC

AD A 048061



# OFFICE OF NAVAL RESEARCH

BRANCH  
OFFICE  
LONDON  
ENGLAND

THIS REPORT IS ISSUED  
FOR INFORMATION PURPOSES  
ON THE UNDERSTANDING  
THAT IT IS NOT A PART OF  
THE SCIENTIFIC LITERATURE  
AND WILL NOT BE CITED  
ABSTRACTED OR REPRINTED

## ONR LONDON CONFERENCE REPORT


C-15-77

ACOUSTIC DETECTION OF NEUTRINO INTERACTIONS IN THE  
OCEAN: THE 1977 DUMAND SUMMER WORKSHOP, MOSCOW,  
26-28 JUNE 1977

A. ROBERTS\*

2 NOVEMBER 1977

\*Fermi Laboratory, Batavia, Illinois



UNITED STATES OF AMERICA

THIS DOCUMENT HAS BEEN APPROVED FOR PUBLIC RELEASE AND SALE; ITS DISTRIBUTION IS UNLIMITED.

AD No. 1

DDC FILE COPY

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

14 REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
ONRL-C-15-77		
4. TITLE (and Subtitle)	5. TYPE OF REPORT & PERIOD COVERED	
Acoustic Detection of Neutrino Interactions in the Ocean: The 1977 <u>DUMAND</u> Summer Workshop, Moscow, 26-28 June 1977.	Conference rept.	
6. AUTHOR(s)	6. PERFORMING ORG. REPORT NUMBER	
A./Roberts, Fermi Laboratory, Batavia, Illinois 60510		
7. PERFORMING ORGANIZATION NAME AND ADDRESS	8. CONTRACT OR GRANT NUMBER(s)	
Office of Naval Research, Branch Office, London Box 39 FPO New York 09510		
9. CONTROLLING OFFICE NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
11. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE	
	2 Nov 1977	
	11	
	15. SECURITY CLASS. (of this report)	
	UNCLASSIFIED	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
	12/14 p.	
16. DISTRIBUTION STATEMENT (of this Report)		
APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
NEUTRINOS                      OPTICAL DETECTORS MUONS                          ACOUSTICAL DETECTION COSMIC RAYS                  PROTON BEAMS ASTROPHYSICS		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
This was the third in a series of Workshops to foster the collaboration of high-energy, cosmic-ray, and theoretical physicists, astrophysicists, astronomers, acousticians, computer scientists, geophysicists, oceanographers, ocean engineers, and other assorted enthusiasts, all captivated by the objective to use the ocean as a gigantic neutrino detector.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

REPORT DOCUMENTATION PAGE	
1. REPORT NUMBER	
2. AUTHOR	
3. PERIODICITY	
4. TITLE	
5. AUTHORING ORGANIZATION	
6. PERFORMING ORGANIZATION	
7. DISTRIBUTION STATEMENT	
8. SECURITY CLASSIFICATION	
9. ABSTRACT	
10. SUBJECT TERMS	
11. DISTRIBUTION STATEMENT	
12. SECURITY CLASSIFICATION	
13. ABSTRACT	
14. SUBJECT TERMS	
15. DISTRIBUTION STATEMENT	
16. SECURITY CLASSIFICATION	
17. ABSTRACT	
18. SUBJECT TERMS	
19. DISTRIBUTION STATEMENT	
20. SECURITY CLASSIFICATION	
21. ABSTRACT	
22. SUBJECT TERMS	
23. DISTRIBUTION STATEMENT	
24. SECURITY CLASSIFICATION	
25. ABSTRACT	
26. SUBJECT TERMS	
27. DISTRIBUTION STATEMENT	
28. SECURITY CLASSIFICATION	
29. ABSTRACT	
30. SUBJECT TERMS	
31. DISTRIBUTION STATEMENT	
32. SECURITY CLASSIFICATION	
33. ABSTRACT	
34. SUBJECT TERMS	
35. DISTRIBUTION STATEMENT	
36. SECURITY CLASSIFICATION	
37. ABSTRACT	
38. SUBJECT TERMS	
39. DISTRIBUTION STATEMENT	
40. SECURITY CLASSIFICATION	
41. ABSTRACT	
42. SUBJECT TERMS	
43. DISTRIBUTION STATEMENT	
44. SECURITY CLASSIFICATION	
45. ABSTRACT	
46. SUBJECT TERMS	
47. DISTRIBUTION STATEMENT	
48. SECURITY CLASSIFICATION	
49. ABSTRACT	
50. SUBJECT TERMS	
51. DISTRIBUTION STATEMENT	
52. SECURITY CLASSIFICATION	
53. ABSTRACT	
54. SUBJECT TERMS	
55. DISTRIBUTION STATEMENT	
56. SECURITY CLASSIFICATION	
57. ABSTRACT	
58. SUBJECT TERMS	
59. DISTRIBUTION STATEMENT	
60. SECURITY CLASSIFICATION	
61. ABSTRACT	
62. SUBJECT TERMS	
63. DISTRIBUTION STATEMENT	
64. SECURITY CLASSIFICATION	
65. ABSTRACT	
66. SUBJECT TERMS	
67. DISTRIBUTION STATEMENT	
68. SECURITY CLASSIFICATION	
69. ABSTRACT	
70. SUBJECT TERMS	
71. DISTRIBUTION STATEMENT	
72. SECURITY CLASSIFICATION	
73. ABSTRACT	
74. SUBJECT TERMS	
75. DISTRIBUTION STATEMENT	
76. SECURITY CLASSIFICATION	
77. ABSTRACT	
78. SUBJECT TERMS	
79. DISTRIBUTION STATEMENT	
80. SECURITY CLASSIFICATION	
81. ABSTRACT	
82. SUBJECT TERMS	
83. DISTRIBUTION STATEMENT	
84. SECURITY CLASSIFICATION	
85. ABSTRACT	
86. SUBJECT TERMS	
87. DISTRIBUTION STATEMENT	
88. SECURITY CLASSIFICATION	
89. ABSTRACT	
90. SUBJECT TERMS	
91. DISTRIBUTION STATEMENT	
92. SECURITY CLASSIFICATION	
93. ABSTRACT	
94. SUBJECT TERMS	
95. DISTRIBUTION STATEMENT	
96. SECURITY CLASSIFICATION	
97. ABSTRACT	
98. SUBJECT TERMS	
99. DISTRIBUTION STATEMENT	
100. SECURITY CLASSIFICATION	

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

ACOUSTIC DETECTION OF NEUTRINO INTERACTIONS  
IN THE OCEAN: THE 1977 DUMAND SUMMER WORKSHOP,  
MOSCOW, 26-28 JUNE 1977

The 1977 Deep Underwater Muon and Neutrino Detector (DUMAND) Summer Workshop was held under the sponsorship and at the invitation of the Soviet Academy of Sciences. Directly following the Neutrino-77 conference at Elbrus which ended 25 June, the Workshop started on a Sunday to avoid delaying the participants who attended both meetings.

The DUMAND project is not yet an officially established one; it has not as yet the blessing of a funding agency, nor does it as yet boast an annual budget. It is a collaboration of high-energy, cosmic-ray, and theoretical physicists, astrophysicists, astronomers, acousticians, computer scientists, geophysicists, oceanographers, ocean engineers, and other assorted enthusiasts, all captivated by its simple but audacious objective: to use the ocean as a gigantic neutrino detector. The aims are to observe and study cosmic-ray neutrinos (and muons) at energies far above those available from any present or foreseeable accelerator—those neutrinos (and muons) that arise from the interactions of the primary cosmic-ray protons (and heavier nuclei) in the earth's atmosphere, as well as those that impinge upon us directly from extraterrestrial sources. The latter have hitherto not been observed. The detection of the low-energy neutrino bursts that accompany gravitational stellar collapse is also one of its aims; but that has been temporarily shelved until someone devises a less-expensive detection procedure for neutrinos in the 20-MeV region than any now known to us.

Such studies would extend our knowledge of the "weak interaction," the sole mechanism of interaction between neutrinos and the rest of the universe, and they would vastly extend the infant science of neutrino astronomy. One might hope to detect neutrino fluxes from the galactic center, from point sources like expanding supernova shells, and perhaps from remote galaxies whose neutrinos date from an early epoch in which the sources may have been much brighter.

Why the ocean? Because detectors of enormous mass are needed to detect the feeble current of neutrinos, which interact with matter so weakly that on the average they may, if their energies are in the right range, readily penetrate the earth, or even a star, without undergoing a single interaction. The ocean alone offers the requisite mass. Two modes of detection appear possible. The more certain, but probably the more expensive, is optical detection: photomultipliers pick up the Cerenkov light that is generated in transparent media by highly relativistic particles—in this case the secondaries produced by the neutrino interaction. This is a standard laboratory method and is well developed and well understood.

ACCESSION FOR	
NTIS	Wire Section <input checked="" type="checkbox"/>
DDC	B.I. Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
CLASSIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
OFFICIAL	
A	

Less certain, but possibly less expensive, is the acoustic method, in which the sound produced by the neutrino interaction is picked up by an array of sensitive hydrophones. The sound results from the pressure pulse generated when the energy of the neutrino, divided among many secondaries, is converted into heat as the particles are brought to rest. This method, first suggested by G. Askarian (Lebedev Inst., Moscow) in the fifties<sup>1</sup> has been shown to work in the laboratory<sup>2,3</sup>, but further studies are still necessary to ascertain how well it will work in the ocean at the energies of interest. Its principal advantage is that sound is attenuated in the ocean far less rapidly than light; thus the detectors may be further apart and fewer in number. Early work on these problems is summarized in the *Proceedings of the DUMAND 1976 Summer Workshop*, published by DUMAND, Fermilab, Batavia, Illinois 60510.

Atmospherically generated neutrinos are accompanied by highly penetrating charged muons, whose intensity at sea level is relatively high. To avoid the interference such particles would produce, it is necessary to go to a considerable depth in the ocean; about 5 to 6 km is desirable. Thus the problem for DUMAND engineers is to design an experimental apparatus in which thousands of sensors, distributed through a volume of perhaps 1 to 100 km<sup>3</sup> of ocean, can be accurately located with respect to each other, and will operate continuously, reliably, and smoothly for many years, returning their data to shore by submarine cable as they are accumulated.

The venue of the Moscow conference emphasized the serious and sustained interest of the Soviet scientists, which has been manifest from the beginning of the project about four years ago. In the last year, following the 1976 DUMAND workshop in Honolulu, experimental and organizational progress has been rapid, and the 1977 conference provided an opportunity for meeting to discuss progress and formulate plans. The DUMAND meeting was held at the Lebedev Institute in Moscow; the arrangements were supervised by Prof. G.T. Zatsepin, and the organization of the conference was in the capable hands of Dr. V.S. Berezinsky (both of the Lebedev Institute).

Two previous workshops, in 1975 and 1976, had been concerned with establishing the conceptual feasibility of both the physical measurements and the ocean engineering. With last year's suggestion of acoustic detection, the economics of the project began to look far more favorable, and the notion of detectors of 10<sup>3</sup> tons (1 km<sup>3</sup>) or even more, began to appear realistic, and even conservative. This year's workshop was concerned with establishing more clearly the possible extraterrestrial sources of ultra-high-energy (UHE) neutrinos and their probable intensities; and in reviewing the progress to date on acoustic detection, both in the USSR and in the US. Total attendance was approximately 25-30, of which eight were US physicists. Soviet attendees included A.I. Alikhanian, B. Pontecorvo, A.E. Chudakov, V.S. Berezinski, G.T.

Zatsepin, L.M. Ozernoi, B.L. Joffe, G. Askarian, B. Dolgoshoin, A.I. Petrukhin, and V.D. Volovik, the latter reporting on his pioneering work on the acoustic detection of particle beams.

The agenda of the three-day meeting, which was conducted on an informal basis (and mostly in English), was varied to suit the interests and needs of the participants, and to follow the directions in which the frequently animated discussions led. On the first day, Prof. F. Reines (Univ. of California, Irvine) described the history of the DUMAND project and its present status; Prof. G.T. Zatsepin (Lebedev Inst. and Univ. of Moscow) reviewed the cosmic-ray properties that determined the conditions for the proposed detection schemes, and discussed what might be learned; and Prof. D.N. Schramm (Univ. of Chicago) discussed the astrophysical aspects, particularly with regard to estimating the anticipated high-energy extraterrestrial neutrino flux.

The uncertainties in these estimates amount to several orders of magnitude, and so predictions range from highly pessimistic (no observable signal) to highly optimistic (many high-energy events).

On the second morning I opened the program with a description of the design of a four-stage prototype  $10^9$ -ton combined optical and acoustic detector. This description elicited the objection that the acoustic threshold of detection (which the Soviet scientists place at about  $10^{15}$  eV, at least ten times the value favored in the US) would be too high for the acoustic detectors to be useful on so small a detector; the event rates would be too low. This was followed by discussions by Ozernoi and by Berezhinski on the probable sources and intensities of extraterrestrial neutrinos. Berezhinsky concluded that with a detector of  $3 \times 10^{11}$  tons ( $300 \text{ km}^3$ ) even a pessimistic estimate of the intensity of neutrinos of  $10^{15}$  eV and above leads to an observed event rate of 10/yr or more.

Prof. L. Sulak (Harvard) discussed the optical detection of high-energy neutrino interactions and the instrumentation required to extract maximum information. The content of his paper was taken mainly from the report of the Neutrino Signature group in the 1976 Summer Study<sup>4</sup>. A purely optical detector is in principle capable of measuring the Bjorken scaling variables  $x$  and  $y$ , by determining the energy of the muon as well as that of the shower; of good accuracy in measuring directions, and moderate accuracy in energy (25%). It might be capable of distinguishing nuclear cascades from electromagnetic showers as well. Given all these data, one can certainly do a cascade measurement that compares favorably with those now customary in accelerator-produced neutrinos, the major difference lying in the inability to determine the signs and energies of the outgoing muons.

This report was received with some skepticism, both as to whether the measurements could indeed be made with the prescribed accuracy, and as to the possibility of achieving this accuracy at a tolerable cost; the system implies an array of optical detectors in the ocean filling a cubic kilometer in a lattice of 40- to 80-m separation.

Berezinsky then discussed experiments<sup>5</sup> to look for resonant production of W bosons (intermediate-vector bosons postulated as the carrier of the weak interaction). The reaction would be  $\bar{\nu}_e + e^- \rightarrow W^- \rightarrow \text{anything}$ . The expected mass of the W is about 70 GeV in the Weinberg form of gauge theory, but they might be much heavier. Ioffe commented that the experiment might possibly determine the mass; he also thought it important (as suggested also by D. Cline) to look for the Higgs boson, a postulated scalar boson necessary in gauge theory.

P. Kotzer (Seattle) gave a short description of a proposed experiment to attempt to observe neutrino oscillations, using an optical detector in the ocean near Seattle, and a neutrino beam produced at Fermilab pointing in that direction. Neutrino oscillations are fluctuations in intensity that might be observed if a coherent beam contains two different types of neutrino whose rest masses are slightly different, and which can transform into each other.

Acoustic Detection of Nuclear Cascades. The third and last day of the meeting was devoted mainly to discussions of acoustic-detection possibilities. First B. Dolgoshein (Moscow Physical Engineering Institute) gave a review of the subject and detailed the results of new Monte Carlo calculations, using the MARS program at Serpukhov<sup>6</sup> to calculate the shower distribution from very high-energy hadronic cascades, and the resultant acoustic radiation pattern (see Fig. 1). As we had earlier speculated, the acoustic pattern is not symmetrical about the plane through the maximum of the cascade and normal to the shower axis. It rises more sharply on the side toward the origin (the upstream side) and falls off more slowly on the downstream side. This implies that with sufficiently accurate observation of the acoustic pattern, the sense of motion of the incident particle along the cascade axis can be inferred. Other new results include the prediction of a considerable acoustic radiation flux in the 100-200 kHz region. The direction of the shower can readily be determined to 10 mrad or better.

V.D. Volovik (Univ. of Kharkov) described the acoustic research experiments that have been going on for several years<sup>7</sup> (since 1971), and which, although published, had been unknown to the DUMAND group. He finds that the mechanism of sound production by particle beams is not purely thermoacoustic, but includes bubble or microbubble formation. The theoretical analysis predicts that as the pressure in the liquid increases beyond 200 atm, the bubble formation contribution vanishes; at the pressures to be encountered in DUMAND (500-600 atm) the effect

becomes pure thermoacoustic, with the exception that dissolved gases, if present, can contribute too.

L. Sulak (Harvard) then presented the results of DUMAND experiments at Brookhaven and Harvard, using proton beams at 150-200 MeV and at 23 GeV. The work of the DUMAND group, which started last October following the suggestion<sup>8</sup> of acoustic detection by Bowen and Dolgoshein at the 1976 Summer Workshop, has now produced significant data<sup>9,10</sup>. The existence of the acoustic signal was readily established. [It had been known for years by Brookhaven technicians, who observed (without amplification) an audible chirp when the 200-MeV linac proton beam was dumped in a large tank of water.] Further efforts were directed at trying to establish the nature of the signal and its threshold of observability. Other experiments concerned the effect of pressure and temperature in the water, dissolved salt, and the question whether the signal was purely thermoacoustic, or whether microbubbles are involved. In the latter case, one would expect different pressure and temperature dependence of the signal. In water, a thermoacoustic signal would be proportional to the parameter  $\beta/C_p$ , where  $\beta$  is the bulk coefficient of thermal expansion and  $C_p$  the specific heat at constant pressure. This quantity is particularly small for water, in which the coefficient of expansion changes sign at 4°C (for fresh water, not for sea water) and for which  $C_p$  is particularly large.

Volovik's experiments, as he reported, showed no vanishing of the signal in water at 4°C, so he concluded that a mechanism other than thermoacoustic must be responsible. The data reported by Sulak did not agree with Volovik's but did not permit unambiguous interpretation.

Dr. John Learned (Univ. of California, Irvine) presented a new exposition of the theory of thermoacoustic pressure pulses produced by particle beams. It uses a Fourier-transform approach in the time domain. It will shortly be published<sup>12</sup>. Pending this, readers may be interested in a simple and readily understood analysis of the acoustic-effect ionizing particles, by Prof. T. Bowen (Univ. of Arizona). It is given in the appendix<sup>13</sup>.

Prof. A. Parvulescu (Institute of Geophysics, Univ. of Hawaii) discussed a few observations on an alternative site for DUMAND, to the west of the Island of Hawaii; and presented some actual spectral analyses of hydrophone recordings made at Barking Sands, Kauai, to show the existence of signals, probably due to porpoises, whose waveforms are remarkably similar to those to be expected from a nuclear cascade.

The meeting closed with expressions of thanks to the organizers, and of hopes for continued and closer cooperation for the future. There are no plans for publishing the proceedings of the workshop.

C-15-77




Fig. 1. Acoustic radiation from a neutrino-produced high-energy cascade. The sonic radiation, being produced in phase along a 10-m path only a few cm in diameter, is emitted in a very narrow-angle disc coaxial with the cascade. The total cascade length is ca. 10 m; the width between half-power points is about 5 m, as shown.

## APPENDIX

THEORY OF ACOUSTIC RADIATION FROM HIGH-ENERGY  
NUCLEAR CASCADES

The theoretical description of the sound waves from particle energy loss has been extended since the Hawaii Workshop by DUMAND participants T. Bowen and J. Learned and also by the Soviet members G.A. Askarian and B.A. Dolgoshein<sup>6</sup>. Bowen and Learned both account<sup>12,13</sup> for the bipolar pulse shapes observed in the Brookhaven experiments and give definite predictions assuming that the spatial distribution of heat deposition is known and that other mechanisms, such as bubble formation, can be neglected. The experimental and theoretical work indicates that particle cascades in water with energies down to  $10^{14} - 10^{15}$  eV should be detectable at one meter by single hydrophones. The task remains to acquire acoustic data from typical high-energy particle cascades both for detailed comparison with theory and for direct use in the design of underwater detectors.

Factors of  $10^3$  or more in sensitivity can be achieved by combining the signals from many hydrophones, so that the detection at distances up to 1 km of cascades of  $10^{14}$  eV will certainly be possible, at the expense of some technical complexity and cost; how much is not yet clear.

The following theoretical analysis is that of T. Bowen<sup>13</sup>.

The acoustic radiation from high-energy cascades in liquids such as water is a theoretical problem considerably simpler than the case of solids, in which shear waves and other complications arise. In liquids the pressure is a scalar quantity, and its propagation is described by the inhomogeneous wave equation

$$\Delta^2 \phi - \frac{1}{c^2} \frac{\partial^2 \phi}{\partial t^2} = - \frac{\beta_0 e_0(\vec{r})}{\rho_0 c_p} \quad (1)$$

where  $\phi$  is a "displacement potential" defined by the equations:

$$q = -\Delta \phi \quad (2)$$

$$p = \rho_0 \partial^2 \phi / \partial t^2 \quad (3)$$

and  $q$  is the displacement of a volume element due to the heat pulse;  
 $p$  is the pressure generated (the acoustic pressure), and

$c$  = velocity of sound in the medium

$\rho_0$  = density of the medium,

$\beta_0$  = coefficient of thermal expansion of the medium


$C_p$  = specific heat at constant pressure and

$e_0(\vec{r})$  = heat deposited adiabatically at  $T = 0$  per unit volume at  $\vec{r}$ .


The solution to this equation is of the form (see, e.g., Morse and Feshbach)

$$\phi(r, t) = \frac{\beta_0}{4\pi\rho_0 C_p} \int_{R < ct} dv \frac{e_0(\vec{r})}{R} \quad (4)$$


where  $\phi(r, t)$  is the retarded potential. The form of  $\phi$  is like this:



(a)



and that of  $p$  is the second time derivative, shown in (b).



(b)

Since  $\phi$  is a constant determined only by the initial heat deposition, we should be able to find a function of  $p$  which gives us  $\phi$ , and thus is a constant of the problem. From the relation between  $p$  and  $\phi$  (Eq. 4), this can be obtained by two successive integrations:

$$I_1(t) = \int_0^t p(t') dt' = \rho_0 \int_0^t \frac{\partial^2 \phi}{\partial t^2} dt' = \rho_0 \left[ \left( \frac{\partial \phi}{\partial t} \right)_t - \left( \frac{\partial \phi}{\partial t} \right)_0 \right]$$

=  $\rho_0 \phi(t)$ , since the second term is zero; and

$$I_2(t) = \int_0^t I_1(t') dt' = \rho_0 \int_0^t \frac{\partial \phi}{\partial t} dt = \rho_0 [\phi(t) - \phi(0)]$$

=  $\rho_0 \phi(t)$ , the second term again vanishing;

$$\text{But now } I_2(t \rightarrow \infty) = \int_0^\infty I_1(t) dt = \rho_0 \phi(t \rightarrow \infty)$$

$$= \frac{\beta_0}{4\pi C_p} \int_{\text{heated region}} dV \frac{e_0(\vec{r})}{R}$$

$$\approx \frac{\beta_0 E_0}{4\pi C_p} \cdot \frac{1}{R} \quad (5)$$

where  $E_0$  is the total energy deposited, provided  $R$  is much larger than the dimensions of the heated region. But now let us evaluate  $I_2(t \rightarrow \infty)$ ; we integrate by parts:

$$\text{Let } u = I_1(t) = \int_0^t p(t') dt'$$

$$du = p(t) dt$$

$$v = t$$

$$dv = dt$$

$$I_2(t \rightarrow \infty) = \left[ t \int_0^t p(t') dt' \right]_0^\infty - \int_0^\infty t p(t) dt$$

The first term vanishes, since there can be no net change in pressure after the event is over; thus

$$\int_0^\infty p(t) dt = - \frac{\beta_0}{4\pi C_p} \int_{\text{heated region}} dv_d \frac{e_0(\vec{r})}{R} \approx \frac{\beta_0 E_0}{4\pi C_p R} \quad (6)$$

The above treatment implies that a simple integration of the pressure pulse in the time domain will give a constant result, characteristic both of the total energy deposited,  $E$ , and of the medium, through the parameter  $\beta/C_p$ . This emphasizes an important feature of this treatment: the mechanism of the signal is assumed to be purely thermoacoustic, with no contributions from microbubbles (as suggested, e.g., by Askarian and Dolgoshein<sup>9</sup>) or ionic expansion (as suggested by Volovik<sup>14</sup>).

In addition, the theory automatically defines the region over which to integrate to get the best signal-to-noise ratio, and also predicts the magnitude of the signal to be observed. A double-integration of the pressure signal, which can be performed with filters, will also give a transformation of the signal into a directly usable output indication of the primary signal strength.

To date, detailed quantitative agreement between theory and observation has not yet been seriously attempted; we believe that the experimentally observed data have uncertainties large enough to cover the discrepancies, which are still of the order of 10 dB. Every effort is being made to diminish these; and since the original discrepancies between theory and experiment were factors of  $10^6$  or more as recently as last fall, we are optimistic about resolving the remaining discrepancy.

## REFERENCES

1. G.A. Askarian, *Atomnaya Energiya* 3, 152 (1957).
2. R.M. White, *J. Appl. Phys.* 34, 3559 (1963).
3. B.L. Beron and R. Hofstadter, *Phys. Rev. Lett.* 23, 184 (1969);  
B.L. Beron *et al*, *IEEE Trans. Nucl. Sci.* 17, 65 (1970).
4. Report of Neutrino Signature Group, P. 297, 1976 SS.
5. V.S. Berezinsky and A.Z. Gazezov, *Letters to JETP* 25, 276 (1977).
6. G.A. Askarian, B.A. Dolgoshein, A.N. Kalinovsky, N.W. Mokhov,  
"Acoustic Detection of High-Energy Neutrinos." P.N. Lebedev Inst.  
Preprint No. 140, Moscow, 1977.
7. V.D. Volovik, V.T. Lazuryk-El'tsufin, *Fiz. Tverd. Tela* 15, 2305  
(1973) in Russian; I.A. Borshkovsky, V.V. Petrenko, V.D. Volovik,  
L.L. Goldin, Ya. L. Kleibock, M.F. Lomanov, *Lett. Nuovo Cim.* 12,  
638 (1975); V.D. Volovik and S.I. Ivanov, *Sov. J. Techn. Phys.*  
20, 1140 (1976) in English translation.
8. T. Bowen, *Proceedings of the DUMAND 1976 Summer Workshop, Honolulu*,  
p. 523. Fermilab, Batavia, Ill., 1976, hereafter referred to as  
1976 SS. G. Askarian and B.A. Dolgoshein, 1976 SS, p. 553; T.  
Bowen, H. Bradner, W.V. Jones, J. Learned, M. Levi, I. Linscott,  
A. Parvulescu, B. Pifer, P. Polakos, J. Strait and L.R. Sulak,  
1976 SS, p. 559.
9. L.R. Sulak, T. Bowen, B. Pifer, P. Polakos, H. Bradner, W.V. Jones,  
J. Learned, T. Armstrong, M. Bregman, M. Levi, J. Strait, I. Linscott,  
A. Parvulescu; "Experimental Studies of the Acoustic Detection  
of Particle Showers and Neutrino Physics Beyond 10 TeV." Presented  
at XIIth Rencontre de Moriond, Flaine, France, Mar. 1977; to be  
published. G.A. Askarian and B.A. Dolgoshein, Lebedev Institute  
Preprint No. 160, 1976. G.A. Askarian, B.A. Dolgoshein, A.N.  
Kalinovsky and N.V. Mokhov, Neutrino '77 Conference, Elbrus,  
Caucasus. V.D. Volovik, V.E. Kobeskoy, IXth All-Union Acoustic  
Conference, paper G3b-10; V.D. Volovik, G.F. Popov, *ibid*, paper  
no. N3G-3; also *Lett. J. Tech. Phys.* 1, 605 (1975).
10. T. Bowen, A. Pifer, P. Polakos, H. Bradner, W.V. Jones, J. Learned,  
T. Armstrong, M. Bregman, M. Levi, J. Strait, L. Sulak, I. Linscott,  
and A. Parvulescu, *Proc. 1977 Nucl. Sci. Symposium*, October, 1977.

11. L. Sulak, unpublished.
12. John G. Learned, Acoustic Radiation by Charged Atomic Particles in Water, An Analysis. To be published.
13. F. Bowen, 15th Int. Conf. on Cosmic Rays, Plovdiv, Bulgaria, Au Aug. 1977.